

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
All ages	7.5	7.5	8.9	2.3	2.1	3.9
Under 65	7.0	7.1	7.4	1.8	1.7	2.7
65 and over	10.8	9.6	19.1	5.7	5.0	12.2
All ages (IARC world std) ^c	5.9	5.9	6.7	1.7	1.6	2.8
<u>Age-Specific Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	-	-	-	-	-
20-24	1.2	1.2	1.2	0.1	0.1	0.2
25-29	4.9	5.4	4.5	0.6	0.6	0.7
30-34	9.7	10.5	8.9	1.3	1.3	1.8
35-39	13.3	14.3	11.4	2.2	2.2	3.1
40-44	14.6	15.1	14.5	3.1	3.0	4.1
45-49	13.2	13.1	14.3	4.1	3.8	6.3
50-54	12.3	11.9	14.9	4.4	4.1	7.0
55-59	11.6	11.2	15.0	4.7	4.3	7.4
60-64	12.1	11.4	16.5	5.2	4.8	8.8
65-69	11.9	11.3	16.1	5.2	4.6	10.1
70-74	11.8	10.3	18.8	5.5	4.8	10.9
75-79	10.0	9.0	18.4	5.7	5.0	12.5
80-84	9.6	8.0	21.2	6.3	5.4	15.1
85+	9.3	7.5	24.8	6.7	5.8	15.4

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.